

< Back to results | 1 of 1

↗ Export

⬇ Download

🖨 Print

✉ E-mail

💾 Save to PDF

★ Add to List

⋮ More... >

Journal of Global Pharma Technology

Volume 10, Issue 8, 2018, Pages 270-275

Alkaloids from Alphonsea elliptica barks and their biological activities

(Article)

Aldulaimi, A.K.O.^a, Azziz, S.S.S.A.^a ✉, Bakri, Y.M.^a, Nafiah, M.A.^a, Awang, K.^b, Aowda, S.A.^c, Litaudon, M.^d, Hassan, N.M.^e, Naz, H.^f, Abbas, P.^g, Hashim, Y.Z.H.-Y.^h, Majhool, A.A.^a 👤

^aDepartment of Chemistry, Faculty of Science and Mathematics, Sultan Idris Education University, 35900 Tanjong Malim, Perak, Malaysia

^bDepartment of Chemistry, Faculty of Science, University of Malaya, Kuala Lumpur, 50603, Malaysia

^cDepartment of Chemistry, Faculty of Science, University Of Babylon, Babylon, Iraq

View additional affiliations ▾

Abstract

▾ View references (23)

This study was carried out to investigate the phytochemicals from the barks of Alphonsea elliptica. The barks were harvested from Hutan Simpan Sembarong, Kluang, Johor; and extracted using dichloromethane (DCM), hexane and methanol successively. Isolation of dichloromethane crude extract was then performed using silica gel column chromatography and preparative thin layer chromatography (PTLC) techniques. Structural identification were carried out via 1D and 2D NMR, UV, IR, MS and comparison with reported data. Phytochemical study of the barks led to the isolation of five alkaloid in which two were oxoaporphines; atherospermidine (1) and liriodenine (2), two were dioxoaporphines; cepharadione A (3) and N-methylouregidione (4), and an azafluorenone alkaloid; kinabaline (5). The hexane, DCM, and methanol crude extracts, together with five isolated alkaloids, were tested for their radical scavenging activity, inhibition of xanthine oxidase activities and cytotoxicity. Atherospermidine and liriodenine possessed antioxidative activities with IC₅₀ value of 20.17 and 10.73, respectively. Atherospermidine, liriodenine, N-methylouregidione and kinabaline showed xanthine oxidase inhibitory activity of 46.29, 7.66, 42.10 and 50.72 μM, respectively. Meanwhile, atherospermidine, liriodenine, cepharadione A and kinabaline showed cytotoxicity against breast cancer cell line with IC₅₀ 89, 86, 79.85 and 62 μg/mL respectively. All the alkaloids tested in this study were isolated for the first time from this species while Cepharadione A is reported for the first time in a species belongs to Alphonsea genus. ©2009-2018, JGPT. All Rights Reserved

SciVal Topic Prominence ⓘ

Topic: Annonaceae | Alkaloids | chemical composition

Prominence percentile: 79.413 ⓘ

Author keywords

Alkaloids | Alphonsea elliptica Hk. f. et Th | Cytotoxicity | Radical scavenging activity | Xanthine oxidase

ISSN: 09758542

Source Type: Journal

Original language: English

Document Type: Article

Publisher: Journal of Global Pharma Technology

References (23)

View in search results format >

Metrics ⓘ View all metrics >

1 Citation in Scopus

0 Field-Weighted Citation Impact



PlumX Metrics ▾

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 1 document

Two New isoquinoline alkaloids from the bark of Alphonsea cylindrica King and their antioxidant activity

Obaid Aldulaimi, A.K. , Abdul Azziz, S.S.S. , Bakri, Y.M. (2019) *Phytochemistry Letters*

View details of this citation

Inform me when this document is cited in Scopus:

Set citation alert >

Set citation feed >

Related documents

A mini review on Alphonsea sp. (Annonaceae): Traditional uses, biological activities and phytochemistry

Bakri, Y.M. , Talip, M.A. , Azziz, S.S.S.A. (2017) *Journal of Applied Pharmaceutical Science*

Antioxidative and cytotoxic activities of crude and isolated compounds of p. Lateriflora (bl.) king

Obaid, A.K. , Azziz, S.S.S.A. , Bakri, Y.M. (2018) *Journal of Pharmaceutical Sciences and Research*

- ☐ 1 Bakri, Y.M., Talip, M.A., Azziz, S.S.S.A.
 A mini review on Alphonsea sp. (Annonaceae): Traditional uses, biological activities and
 phytochemistry ([Open Access](#))

(2017) *Journal of Applied Pharmaceutical Science*, 7 (10), pp. 200-203. Cited 2 times.
http://www.japsonline.com/admin/php/uploads/2463_pdf.pdf
 doi: 10.7324/JAPS.2017.71030

[View at Publisher](#)

- ☐ 2 e Silva, M.G., De Oliveira, A.P., Araújo, C.S., De Lavor, É.M., Silva, J.C., Mendes, R.L., Pessoa, C.Ó., (...), Almeida, J.R.S.
 Phytochemical screening, cytotoxicity and acute toxicity of Annona vepretorum Mart
 (Annonaceae) leaf extracts ([Open Access](#))

(2017) *Tropical Journal of Pharmaceutical Research*, 16 (3), pp. 597-604. Cited 2 times.
https://www.tjpr.org/admin/12389900798187/2017_16_3_14.pdf
 doi: 10.4314/tjpr.v16i3.14

[View at Publisher](#)

- ☐ 3 González-Esquinca, A.R., De-La-Cruz-Chacón, I., Castro-Moreno, M., Orozco-Castillo, J.A., Riley-Saldaña, C.A.
 Alkaloids and acetogenins in annonaceae development: Biological considerations ([Open Access](#))

(2014) *Revista Brasileira de Fruticultura*, 36 (SPEC. EDITION 1), pp. 1-16. Cited 16 times.
<http://www.scielo.br/pdf/rbf/v36nspe1/v36nspe1a01.pdf>
 doi: 10.1590/S0100-29452014000500001

[View at Publisher](#)

- ☐ 4 Husain, K., Jamal, J.A., Jalil, J.
 Phytochemical study of Cananga odorata Hook, F. & Thoms (Annonaceae)

(2012) *International Journal of Pharmacy and Pharmaceutical Sciences*, 4 (SUPPL. 4), pp. 465-467. Cited 6 times.
<http://www.ijppsjournal.com/Vol4Suppl4/4545.pdf>

- ☐ 5 Turner, I.M., Utteridge, T.M.A.
 A new species of alphonsea (Annonaceae) from Peninsular Malaysia

(2015) *Blumea: Journal of Plant Taxonomy and Plant Geography*, 59 (3), pp. 206-208. Cited 6 times.
<http://www.ingentaconnect.com/search/download?pub=infobike%3a%2f%2fnhn%2fblumea%2f2015%2f00000059%2f00000003%2fart00007&mimetype=application%2fpdf>
 doi: 10.3767/000651915X688443

[View at Publisher](#)

- ☐ 6 (2013) *Version 1.1*. Cited 73 times.
 Published on the Internet; accessed 1st
[The Plant List](#)

- ☐ 7 Turner, I.M.
 Notes on the Annonaceae of the Malay Peninsula
 (2016) *Gard. Bull. Singapore*, 68 (1), p. 65. Cited 5 times.